

ANOSEA

MECLIZINE USP

Compositions:

Each film-coated tablet contains Meclizine HCl USP 50 mg.

Pharmacology:

Anosea (Meclizine HCl) is a first generation piperazine-derivative antihistamine that shows marked protective activity against nebulized histamine and used as an antiemetic.

Dosage And Administration:

The usual doses of Anosea (Meclizine HCl) are as follow Nausea and vomiting: 25-50 mg once daily or as directed by physician. Motion sickness: 25-50 mg once daily (1/2 or 1 tablets 1 hour before to travel starts, The dose may be repeated every 24 hours during the journey. Vertigo: 25-100mg daily in divided doses. Radiotherapy induced Nausea and vomiting: 50 mg (Meclizine Hydrochloride) administered 2 to 12 hours prior to radiotherapy.

Contraindications:

Meclizine HCl is contraindicated who have shown hypersensitivity to it.

Warning And Precaution:

Patients with asthma, bronchitis, emphysema, enlarged prostate, glaucoma or urinary tract blockage should take Meclizine (like other antiemetics) with caution.

Side Effects:

Common side effects are Drowsiness, dry mouth, and on rare occasions, blurred vision have been reported.

Use in Pregnancy and Lactation:

Meclizine shows no risk of birth defects or abnormalities when administered during pregnancy. Although Meclizine may excrete into the milk, it causes no harm in nursing babies.

Drug Interaction:

Meclizine may increase the toxicity with CNS depressants, neuroleptics and anticholinergic. Alcohol, sedatives and tranquilizers can increase the drowsiness of the patient

Overdosage:

Symptoms: Extreme excitability, seizures, drowsiness and hallucinations. Treatment: Appropriate supportive and symptomatic treatment. Consider dialysis.

Storage:

Store in a cool (below 30°C) and dry place. Keep away from light and out of reach of children.

Packing:

Each box contains 5x10's and 10x10's tablet in blister pack.

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.

Shafipur, Gazipur, Bangladesh.