

SERTAGIN

SERTACONAZOLE BP

Compositions:

Cream: Each gm cream contains Sertaconazole nitrate BP 20 mg.

Pharmacology:

Sertaconazole, an imidazole antifungal agent, inhibits fungal cytochrome P450-mediated 14 alpha-lanosteroldemethylase enzyme. Ergosterol is the key component of fungal cell membranes and lack of this component leads to fungal cell injury by leakage of key constituents in the cytoplasm from the cell.

Dosage And Administration:

In the treatment of interdigitaltineapedis, Sertaconazole cream should be applied twice daily for 4 weeks. A sufficient amount of Sertaconazole cream should be applied to cover both the affected areas between the toes and the immediately surrounding healthy skin of patients with interdigitaltineapedis. Not for ophthalmic, oral, or intravaginal use.

Contraindications:

undefined

Warning And Precaution:

undefined

Side Effects:

Most common adverse reactions observed in clinical trials are contact dermatitis, dry skin, burning skin, application site skin tenderness.

Use in Pregnancy and Lactation:

The Pregnancy Category C. Sertaconazole is not recommended during pregnancy & lactation

Drug Interaction:

Potential interactions between Sertaconazole cream and other drugs have not been Systematically evaluated

Overdosage:

Over dosage withSertaconazole 2% cream has not been reported to date

Storage:

Store in a cool (below 30°C) and dry place. Keep away from light and out of reach of children.

Packing:

Each lamitube contains 20 gm cream..

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.

Shafipur, Gazipur, Bangladesh.