

VOTAB

VORICONAZOLE USP

Compositions:

Tablet: Each film coated tablet contains Voriconazole USP 200 mg.

Pharmacology:

Voriconazole is a triazole antifungal medication used to treat serious fungal infections. The primary mode of action of Voriconazole is the inhibition of fungal cytochrome P450 mediated 14 alpha-lanosterol demethylation, an essential step in fungal ergosterol biosynthesis

Dosage And Administration:

Voriconazole is to be taken at least one hour before or one hour following a meal. For Patients 40 kg or above: Loading Dose: 400 mg twice daily (for the first 24 hours). Maintenance Dose: 200 mg twice daily (from the 2nd day). For Patients less than 40 kg: Loading Dose: 200 mg twice daily (for the first 24 hours) Maintenance Dose: 100 mg or 150 mg twice daily (from the 2nd day). Treatment duration should be as short as possible depending the patient's clinical and mycological response.

Contraindications:

Known hypersensitivity to Voriconazole or any other components of this drug- • Co-administration with terfenadine, astemizole, cisapride, pimozide or quinidine, sirolimus due to risk of serious adverse reactions • Co-administration with rifampin, carbamazepine, long-acting barbiturates, efavirenz, ritonavir, rifabutin, ergot alkaloids and St. John's Wort due to risk of loss of efficacy

Warning And Precaution:

Long term exposure (treatment or prophylaxis) greater than 180 days requires careful assessment of the benefit-risk balance. Squamous cell carcinoma of the skin (SCC) has been reported in relation with long-term voriconazole treatment.

Side Effects:

The most common side effects are abdominal pain, anemia, blurred vision, headache, chest pain, nausea and diarrhea.

Use in Pregnancy and Lactation:

There are no adequate and well-controlled studies in pregnant woman. It should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. The excretion of Voriconazole into breast milk has not been investigated.

Drug Interaction:

CYP3A4, CYP2C9 and CYP2C19 inhibitors and inducers: Adjust voriconazole dosage and monitor for adverse reactions or lack of efficacy. • Voriconazole may increase the concentrations and activity of drugs that are CYP3A4, CYP2C9 and CYP2C19 substrates. Reduce dosage of these other drugs and monitor for adverse reactions. • Phenytoin or Efavirenz: with co-administration, increase maintenance oral and intravenous dosage of Voriconazole

Overdosage:

In an overdose, haemodialysis may assist in the removal of Voriconazole and SBECD from the body

Storage:

Store in a cool (below 30°C) and dry place. Keep away from light and out of reach of children.

Packing:

Each box contains 1x10 tablets in Alu-Alu blister pack

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.

Shafipur, Gazipur, Bangladesh.