

DEXON G

DEXAMETHASONE BP + GATIFLOXACIN SESQUIHYDRATE INN

Compositions:

Each ml Contains Gatifloxacin Sesquihydrate INN equivalent to Gatifloxacin 3 mg and Dexamethasone Sodium Phosphate BP equivalent to Dexamethasone Phosphate 1mg.

Pharmacology:

Gatifloxacin inhibits bacterial DNA gyrase and Topoisomerase IV, the enzymes responsible for replication, transcription and repair of bacterial DNA. Dexamethasone suppresses the inflammation by decreasing the production of inflammatory mediators through the inhibition of the enzyme Phospholipase A2.

Dosage And Administration:

Instill 1-2 drops into the conjunctival sac 3 to 4 times daily.

Contraindications:

Contraindicated in fungal disease of ocular structures, viral conjunctivitis and in patients who are hypersensitive to any of the components of this preparation.

Warning And Precaution:

For ophthalmic use only. If the product is used for 10 days or more, intraocular pressure should be monitored routinely.

Side Effects:

Mild burning, chemosis, redness may occur.

Use in Pregnancy and Lactation:

Pregnancy: There are no adequate and well-controlled studies in pregnant women. Dexon-G should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Lactation: Caution should be exercised when Dexon-G is administered to a nursing mother.

Drug Interaction:

There are no known drug interactions.

Overdosage:

Overdosage of topical application of acyclovir 3% is unlikely because of minimal absorption.

Storage:

Store in a cool and dry place, away from light. Keep out the reach of children. After one month of the opening do not use the medicine of dropper.

Packing:

Each plastic dropper bottle contains 5 ml of sterile eye drops.

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.

Shafipur, Gazipur, Bangladesh.