

FLOROMOX XG

MOXIFLOXACIN HYDROCHLORIDE BP

Compositions:

Each ml contains Moxifloxacin Hydrochloride BP equivalent to Moxifloxacin 5 mg (enriched with Xanthan Gum BP 0.4%).

Pharmacology:

Moxifloxacin shows antibacterial activity by inhibiting the bacterial enzyme topoisomerase-II (DNA gyrase) and topoisomerase-IV.

Dosage And Administration:

One drop in the affected eye's twice daily for 7 days.

Contraindications:

No information available.

Warning And Precaution:

Prolonged use may result in overgrowth of non-susceptible organisms, including fungi like with other anti-infectives.

Side Effects:

The most frequently reported ocular side effects are ocular discomfort, irritation etc. may occur.

Use in Pregnancy and Lactation:

Pregnancy: Moxifloxacin should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus as there are no adequate and well-documented studies in pregnant women. Lactation: Moxifloxacin has not been measured in human milk, although it can be presumed to be excreted in human milk. Caution should be exercised when Moxifloxacin is administered to a lactating mother.

Drug Interaction:

Drug-drug interaction studies have not been conducted with Moxifloxacin Hydrochloride. In vitro studies indicate that Moxifloxacin does not inhibit CYP3A4, CYP2D6, CYP2C9, CYP2C19 or CYP1A2 indicating that Moxifloxacin is unlikely to alter the pharmacokinetics of drugs metabolized by these cytochrome P450 isoenzymes.

Overdosage:

No information available.

Storage:

Store below 30° C at cool and dry place, protect from light. Keep out of reach of children.

Packing:

Each plastic dropper bottle contains 5 Sterile & Pyrogen free Ophthalmic Solution.

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.

Shafipur, Gazipur, Bangladesh.