

OBACTIN

OFLOXACIN USP

Compositions:

Obactin sterile eye drops: Each ml contains Ofloxacin USP 3 mg.

Pharmacology:

Ofloxacin is a bacterial DNA gyrase inhibitor, the enzyme responsible for duplication, transcription and repair of bacterial DNA.

Dosage And Administration:

Instill 1 drop in the affected eye(s) every 2 to 4 hours for the first two days and then 4 times daily. The length of treatment should not exceed ten days.

Contraindications:

Obactin Eye Drops is contraindicated in patients who are hypersensitive to Ofloxacin or any other component of this preparation.

Warning And Precaution:

Prolonged use of eye drops may result in overgrowth of non-susceptible organisms and secondary infection respectively.

Side Effects:

Transient ocular irritation, burning, stinging, redness, itching or photophobia have been reported.

Use in Pregnancy and Lactation:

There are no adequate and well-controlled studies in pregnant women. This drug should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Use in lactation: It is not known whether Ofloxacin is excreted in human milk. Cautions should be taken when Ofloxacin eye & ear drops is administered during lactation.

Drug Interaction:

No information available.

Overdosage:

No information provided.

Storage:

Store at room temperature (15°-25°C) and dry place protected from light. Do not use more than 30 days after first opening of the bottle. Keep out of the reach of children.

Packing:

Obactin Eye Drops: Each plastic dropper bottle contains 5 ml sterile Eye Drops.

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.

Shafipur, Gazipur, Bangladesh.