

# **OBACTIN**

OFLOXACIN USP

## **Compositions:**

Obactin sterile eye drops: Each ml contains Ofloxacin USP 3 mg.

## **Pharmacology:**

Ofloxacin is a bacterial DNA gyrase inhibitor, the enzyme responsible for duplication, transcription and repair of bacterial DNA.

## **Dosage And Administration:**

Instill 1 drop in the affected eye(s) every 2 to 4 hours for the first two days and then 4 times daily. The length of treatment should not exceed ten days.

## **Contraindications:**

Obactin Eye Drops is contraindicated in patients who are hypersensitive to Ofloxacin or any other component of this preparation.

## **Warning And Precaution:**

Prolonged use of eye drops may result in overgrowth of non-susceptible organisms and secondary infection respectively.

## **Side Effects:**

Transient ocular irritation, burning, stinging, redness, itching or photophobia have been reported.

## **Use in Pregnancy and Lactation:**

There are no adequate and well-controlled studies in pregnant women. This drug should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Use in lactation: It is not known whether Ofloxacin is excreted in human milk. Cautions should be taken when Ofloxacin eye & ear drops is administered during lactation.

## **Drug Interaction:**

No information available.

## **Overdosage:**

No information provided.

## **Storage:**

Store at room temperature (15°-25°C) and dry place protected from light. Do not use more than 30 days after first opening of the bottle. Keep out of the reach of children.

## **Packing:**

Obactin Eye Drops: Each plastic dropper bottle contains 5 ml sterile Eye Drops.

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.

Shafipur, Gazipur, Bangladesh.