

IRYSIN

MEBEVERINE HYDROCHLORIDE BP

Compositions:

Irisyn Tablet: Each film-coated tablet contains Mebeverine Hydrochloride BP 135 mg.

Pharmacology:

Irisyn is a musculotropic antispasmodic agent used to relieve cramps or spasms of the stomach and intestine. It is particularly useful in treating irritable bowel syndrome (IBS) and similar conditions. It works by relaxing the muscles and helping to restore the normal movement of the gut.

Dosage And Administration:

Adults, elderly and children over 10 years : One tablet three times a day, preferably 20 minutes before meal. After a period of several weeks when the desired effect has been obtained, the dosage may be gradually reduced. children: Not recommended for children under 10 years.

Contraindications:

Mebeverine is contraindicated in patients who are hypersensitive to any component of this product.

Warning And Precaution:

Neonates: Not recommended for neonates. Children: The preparation is recommended for children over 10 years.

Side Effects:

Generally Mebeverine is well tolerated. However, few side effects like skin rash, urticaria and angioedema may appear.

Use in Pregnancy and Lactation:

Pregnancy: No teratogenicity has been shown in animal experiments and therefore, there is no known teratogenic effect in human.

Drug Interaction:

Concomitant use of two or more such drugs can increase side effects such as dry mouth, urine retention and constipation.

Overdosage:

If you or someone else take more Mebeverine Tablets than you should (overdose), contact your doctor or hospital emergency department immediately.

Storage:

Store in a cool (below 30°C) and dry place protected from light and moisture. Keep out of the reach of children.

Packing:

Irisyn Tablet: Each box contains 5x10 tablet in blister pack.

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.

Shafipur, Gazipur, Bangladesh.