

AMINOSIN PLUS

AMINO ACID WITH GLUCOSE & ELECTROLYTES

Compositions:

Each 100 ml IV Infusion contains Essential Amino Acids Specification Quantity L-Isoleucine USP 0.390 g L-Leucine USP 0.530 g L-Lysine Hydrochloride USP 0.390 g L-Methionine USP 0.190 g L-Phenylalanine USP 0.550 g L-Threonine USP 0.300 g L-Tryptophan USP 0.100 g L-Valine USP 0.430 g L-Histidine USP 0.240 g L-Tyrosine USP 0.050 g Non-Essential Amino Acids L-Arginine USP 0.330 g L-Aspartic Acid USP 0.410 g L-Glutamic Acid BP 0.900 g L-Alanine USP 0.300 g L-Cysteine BP 0.140 g Glycine USP 0.210 g L-Proline USP 0.810 g L-Serine USP 0.750 g Carbohydrate Anhydrous Glucose BP 10.000 g Electrolytes (mmol/L) Sodium (Na⁺) 50.0 Potassium (K⁺) 20.0 Calcium (Ca⁺⁺) 2.5 Magnesium (Mg⁺⁺) 1.5 Chloride (Cl⁻) 32.0

Pharmacology:

Aminosin Plus is a sterile aqueous solution of Amino Acids IV Infusion and 10% Glucose with electrolytes, which is necessary as the nitrogen sources for parenteral nutrition. Nitrogen is provided in the form of essential and non-essential amino acids.

Indications:

As a source of amino acids, Glucose and electrolytes in adult and pediatric patients needing IV nutrition. Amino Acid IV Infusion with 10% Glucose and Electrolytes is particularly suitable for patients with basal amino acids requirements.

Dosage And Administration:

Adult: The nitrogen requirement for maintenance of body protein mass depends on the condition of the patient (nutritional state and degree of metabolic stress). The requirements are 0.1-0.15 g nitrogen/kg body weight/day (no or minor metabolic stress and normal nutritional state), 0.15-0.2 g nitrogen/kg body weight/day (moderate metabolic stress with or without malnutrition) and up to 0.2-0.25 g nitrogen/kg body weight/day (severe catabolism as in burns, sepsis and trauma). The dosage range 0.1-0.25 g nitrogen/kg body weight/day corresponds to 11-27 mL Amino Acid IV Infusion with 10% Glucose and Electrolytes /kg body weight/day, respectively. In obese patients, the dose should be based on the estimated ideal weight. Depending upon patient requirements, up to 1000-2000 mL of Amino Acid IV Infusion with 10% Glucose and Electrolytes may be infused IV per 24 hrs. Amino Acid IV Infusion with 10% Glucose and Electrolytes should be infused slowly, at a rate not exceeding 1000 mL in 6 hrs corresponding to approximately 2.8 mL/min. In patients with basal amino acids requirements, the less concentrated Amino Acid IV Infusion with 10% Glucose and Electrolytes may be used. Infant and Children: In children and infants, a maximal rate of infusion of 30 mL Amino Acid IV Infusion with 10% Glucose and Electrolytes/kg body weight/day is recommended, with a stepwise increase in the rate of administration during the 1st week of treatment

Contraindications:

Patients with inborn errors of amino acid metabolism, severe liver dysfunction and in severe uremia when dialysis facilities are not available. Due to the content of Glucose, Amino Acids IV Infusion and 10% Glucose with electrolytes is contraindicated in patients with hyperosmolar nonketotic diabetic coma.

Warning And Precaution:

IV infusion of amino acids is accompanied by increased urinary excretion of the trace elements copper and, in particular zinc, which should be taken into account in the dosing of trace elements, particularly during long-term IV nutrition. Hyperphenylalaninemia has been

Shafipur, Gazipur, Bangladesh.