## ANOSEA

#### MECLIZINE USP

### **Compositions:**

Each film-coated tablet contains Meclizine HCl USP 50 mg.

# Pharmacology:

Anosea (Meclizine HCl) is a first generation piperazine-derivative antihistamine that shows marked protective activity against nebulized histamine and used as an antiemetic.

#### **Indications:**

For the prevention and treatment of nausea, vomiting, dizziness, motion sickness and vertigo.

### **Dosage And Administration:**

The usual doses of Anosea (Meclizine HCl) are as follow Nausea and vomiting: 25-50 mg once daily or as directed by physician. Motion sickness: 25-50 mg once daily (1/2 or 1 tablets 1 hour before to travel starts, The dose may be repeated every 24 hours during the journey. Vertigo: 25-100mg daily in divided doses. Radiotherapy induced Nausea and vomiting: 50 mg (Meclizine Hydrochloride) administered 2 to 12 hours prior to radiotherapy.

#### **Contraindications:**

Meclizine HCl is contraindicated who have shown hypersensitivity to it.

#### **Warning And Precaution:**

Patients with asthma, bronchitis, emphysema, enlarged prostate, glaucoma or urinary tract blockage should take Meclizine (like other antiemetics) with caution.

### **Side Effects:**

Common side effects are Drowsiness, dry mouth, and on rare occasions, blurred vision have been reported.

# **Use in Pregnancy and Lactation:**

Meclizine shows no risk of birth defects or abnormalities when administered during pregnancy. Although Meclizine may excrete into the milk, it causes no harm in nursing babies.

#### **Drug Interaction:**

Meclizine may increase the toxicity with CNS depressants, neuroleptics and anticholinergic. Alcohol, sedatives and tranquilizers can increase the drowsiness of the patient

#### Overdosage:

Symptoms: Extreme excitability, seizures, drowsiness and hallucinations. Treatment: Appropriate supportive and symptomatic treatment. Consider dialysis.

### **Storage:**

Store in a cool (below 30°C) and dry place. Keep away from light and out of reach of children.

# Packing:

Each box contains 5x10's and 10x10's tablet in blister pack.

Manufactured By: The IBN SINA Pharmaceutical Industry PLC. Shafipur, Gazipur, Bangladesh.